

11/22/00
JC952 U.S. PTO

11-24-00

A

MAFE
jc915 U.S. PTO
09/72/114
11/22/00

Applicant(s) : Hirohiko Hirochika, Muneo Yamazaki and
Akio Miyao

For : A NOVEL GENE INVOLVED IN BRASSINOSTEROID
RESPONSES


EXPRESS MAIL CERTIFICATION

"Express Mail" mailing label number EI124987820US.

Date of Deposit November 22, 2000.

I hereby certify that this transmittal letter and the other papers and fees identified in this transmittal letter as being transmitted herewith are being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 C.F.R. § 1.10 on the date indicated above and are addressed to the Hon. Commissioner for Patents, Washington, D.C. 20231.

Hon. Commissioner
for Patents
Washington, D.C. 20231
Box PATENT APPLICATION


CATHERINE S. SAMML
New York, New York
November 22, 2000

TRANSMITTAL LETTER FOR
ORIGINAL PATENT APPLICATION

Sir:

Transmitted herewith for filing are the
[X] specification; [X] claims; [X] abstract; [X] declaration
and [X] power of attorney, for the above-identified patent
application.

Also transmitted herewith are:

[X] 7 sheets of:

[X] Formal drawings.

[] Informal drawings. Formal drawings will be filed
during the pendency of this application.

☐ Certified copy(ies) of application(s)

(country)	(appln. no.)	(filed)
(country)	(appln. no.)	(filed)
(country)	(appln. no.)	(filed)

from which priority is claimed.

☐ An assignment of the invention to _____.

☐ A check in the amount of \$40.00 to cover the recording fee.

☐ Please charge \$40.00 to Deposit Account No. 06-1075 in payment of the recording fee. A duplicate copy of this transmittal letter is transmitted herewith.

☐ An associate power of attorney.

The filing fee has been calculated as shown below:

FOR	NUMBER FILED	NUMBER EXTRA	RATE	FEE
BASIC FEE				\$710.00
TOTAL CLAIMS	3	- 20 = 0	X \$ 18 =	\$ 0.00
INDEPENDENT CLAIMS	1	- 3 = 0	X \$ 80 =	\$ 0.00
<input type="checkbox"/> MULTIPLE DEPENDENT CLAIMS			+ \$270 =	\$ 0.00
			TOTAL	<u>\$710.00</u>

☒ A check in the amount of \$710.00 in payment of the filing fee is transmitted herewith.

[] This application is being filed unaccompanied by a filing fee. The appropriate filing fee will be paid in response to a Notice to File Missing Parts, pursuant to 37 C.F.R. § 1.53(f).


[X] The Commissioner is hereby authorized to charge payment of any additional filing fees required under 37 C.F.R. § 1.16, in connection with the paper(s) transmitted herewith, or credit any overpayment of same, to deposit Account No. 06-1075. A duplicate copy of this transmittal letter is transmitted herewith.

[] Amend the specification by inserting before the first line the sentence: -- This is a [] continuation-in-part, of application Serial No.: _____, filed

entitled _____

_____.

[] Please charge \$_____ to Deposit Account No. 06-1075 in payment of the filing fee. A duplicate copy of this transmittal letter is transmitted herewith.



James F. Haley, Jr. (Reg. No. 27,794)
Attorney for Applicants
c/o FISH & NEAVE
Customer No. 1473
1251 Avenue of the Americas
New York, New York 10020-1104
Tel.: (212) 596-9000

INVENTOR INFORMATION

Inventor One Given Name:: Hirohiko
Family Name:: Hirochika
Postal Address Line One:: c/o Ministry Agric., Forestry & Fisheries
Postal Address Line Two:: 2-1-2, Kannondai, Tsukuba-shi
City:: Ibaraki
Country:: Japan
Postal or Zip Code:: 305-8602
Citizenship Country:: Japan
Inventor Two Given Name:: Muneo
Family Name:: Yamazaki
Postal Address Line One:: c/o Ministry Agric., Forestry & Fisheries
Postal Address Line Two:: 2-1-2, Kannondai, Tsukuba-shi
City:: Ibaraki
Country:: Japan
Postal or Zip Code:: 305-8602
Citizenship Country:: Japan
Inventor Three Given Name:: Akio
Family Name:: Miyao
Postal Address Line One:: c/o Ministry Agric., Forestry & Fisheries
Postal Address Line Two:: 2-1-2, Kannondai, Tsukuba-shi
City:: Ibaraki
Country:: Japan
Postal or Zip Code:: 305-8602
Citizenship Country:: Japan

CORRESPONDENCE INFORMATION

Correspondence Customer Number:: 1473
Fax One:: 212-596-9090
Electronic Mail One:: jhaley@fishneave.com

APPLICATION INFORMATION

Title Line One:: A NOVEL GENE INVOLVED IN BRASSINOSTEROID
Title Line Two:: RESPONSES
Total Drawing Sheets:: 7
Formal Drawings?: Yes
Application Type:: Utility
Docket Number:: MAFF-1
Secrecy Order in Parent Appl.?: No

REPRESENTATIVE INFORMATION

Representative Customer Number:: 1473
Registration Number One:: 27794

PRIOR FOREIGN APPLICATIONS

Foreign Application One:: 2000-149106
Filing Date:: 05-19-2000
Country:: Japan
Priority Claimed:: Yes

Source:: PrintEFS Version 1.0.1

AR022

A NOVEL GENE INVOLVED IN BRASSINOSTEROID RESPONSES

EI124987820US

BACKGROUND OF THE INVENTION

1. FIELD OF THE INVENTION:

5 The present invention relates to a novel gene. In particular, the present invention relates to a novel gene in plants which encodes a protein having the function of controlling an in-vivo signal transduction system in a physiological reaction system against brassinosteroid
10 hormone.

2. DESCRIPTION OF THE RELATED ART:

15 Transposons are mutagenic genes which are known to be ubiquitous in animal, yeast, bacterial, and plant genomes. Transposons are classified into two classes, Class I and Class II, depending on their transposition mechanisms. Transposons belonging to Class II are transposed in the form of DNAs without being replicated. Known Class II
20 transposons include the Ac/Ds, Spm/dSpm and Mu elements of Zea mays (Fedoroff, 1989, Cell 56, 181-191; Fedoroff et al., 1983, Cell 35, 235-242; Schiefelbein et al., 1985, Proc. Natl. Acad. Sci. USA 82, 4783-4787), and the Tam element of Antirrhinum majus (Bonas et al., 1984, EMBO J., 3,
25 1015-1019). Class II transposons are widely used for gene isolation techniques which utilize transposon tagging. Such techniques utilize the fact that a transposon induces physiological and morphological changes when inserted into genes. The affected gene can be isolated by detecting such
30 changes (Bancroft et al., 1993, The Plant Cell, 5, 631-638; Colasanti et al., 1998, Cell, 93, 593-603; Gray et al., 1997, Cell, 89, 25-31; Keddie et al., 1998, The Plant Cell, 10, 877-887; Whitham et al., 1994, Cell, 78, 1101-1115).

Transposons belonging to Class I, also referred to as retrotransposons, are replicated and transposed via RNA intermediates. Class I transposons were first identified and characterized in *Drosophila* and in yeasts. However, recent studies have revealed that Class I transposons are ubiquitous in plant genomes and account for a substantial portion of the genomes (Bennetzen, 1996, *Trends Microbiolo.*, 4, 347-353; Voytas, 1996, *Science*, 274, 737-738). A large majority of retrotransposons appear to be inactive. Recent studies indicate that some of these retrotransposons are activated under stress conditions such as injuries, pathogenic attacks, or cell culture (Grandbastien, 1998, *Trends in Plant Science*, 3, 181-187; Wessler, 1996, *Curr. Biol.* 6, 959-961; Wessler et al., 1995, *Curr. Opin. Genet. Devel.* 5, 814-821). Activation under stress conditions has been reported for *Tnt1A* and *Ttol* in tobacco (Pouteau et al., 1994, *Plant J.*, 5, 535-542; Takeda et al., 1988, *Plant Mol. Biol.*, 36, 365-376), and *Tos17* in rice (Hirochika et al., 1996, *Proc. Natl. Acad. Sci. USA*, 93, 7783-7788), for example.

The *Tos17* retrotransposon of rice is one of the most-extensively studied plant Class I elements in plants. *Tos17* was cloned by an RT-PCR method using a degenerate primer prepared based on a conservative amino acid sequence in reverse transcription enzyme domains between *Tyl-copia* retroelements (Hirochika et al., 1992, *Mol. Gen. Genet.*, 233, 209-216). *Tos17* is 4.3kb long, and has two 138bp LTRs (long chain terminal repetitions) and PBS (primer binding sites) complementary to the 3' end of the start methionine tRNA (Hirochika et al., 1996, *supra*). *Tos17* transcription is strongly activated through tissue culture, and its copy

number increases with culture time. In Nipponbare, a model Japonica cultivar used for genome analysis, two copies of Tos17 are initially present, which are increased to 5 to 30 copies in a regenerated plant after tissue culture (Hirochika et al., 1996, supra). Unlike Class II transposons which were characterized in yeasts and Drosophila, Tos17 is transposed in chromosomes in random manners and causes stable mutation, and therefore provides a powerful tool for functional analysis of rice genes (Hirochika, 1997, Plant Mol. Biol. 35, 231-240; 1999, Molecular Biology of Rice (ed. by K. Shimamoto, Springer-Verlag, 43-58).

SUMMARY OF THE INVENTION

The present invention relates to a polynucleotide encoding a plant gene capable of controlling a signal transduction system for brassinosteroid hormone, the polynucleotide encoding an amino acid sequence from Met at position 1 to Arg at position 1057 of SEQ ID NO: 2 in the SEQUENCE LISTING, including any polynucleotide encoding an amino acid sequence in which one or more amino acids are deleted, substituted or added to the amino acid sequence.

In one embodiment of the invention, the polynucleotide may be derived from rice.

In another embodiment of the invention, the polynucleotide may be as represented by SEQ ID NO: 1 in the SEQUENCE LISTING.

The present invention further relates to methods for controlling various effects in plants in which

brassinosteroid hormone is involved, e.g., growth promotion, yield increase, quality improvement, maturation enhancement, and tolerance against biotic and abiotic stresses.

5

The inventors diligently conducted systematic analyses of phenotypes of plants having a newly transposed Tos17 copy and sequences adjoining Tos17 target sites with respect to rice. As a result, the inventors found a dwarf rice mutation obtained from Tos17 insertion, and isolated the gene responsible for this mutation by utilizing Tos17 as a tag, thereby accomplishing the present invention.

10

Thus, the invention described herein makes possible the advantage of providing a novel plant gene which can be provided by using Tos17.

15

This and other advantages of the present invention will become apparent to those skilled in the art upon reading and understanding the following detailed description with reference to the accompanying figures.

20

BRIEF DESCRIPTION OF THE DRAWINGS

25

Figures 1A and 1B are photographs showing a brassinosteroid non-sensitive mutant having Tos17 inserted therein, which was found among regenerated Akitakomachi lineage. On the left of each figure is shown a brassinosteroid non-sensitive mutant having Tos17 inserted therein. On the right of each figure is shown a wild type plant body. Figure 1A evidences an influence toward dwarfism and upright form. Figure 1B evidences an influence toward malformation of grain hulls.

30

Figures 2A and 2B are photographs showing a brassinosteroid non-sensitive mutant having Tos17 inserted therein, which was found among regenerated Nipponbare lineage. On the left of each figure is shown a brassinosteroid non-sensitive mutant having Tos17 inserted therein. On the right of each figure is shown a wild type plant body. Figure 2A evidences an influence toward dwarfism and upright form. Figure 2B evidences an influence toward malformation of grain hulls.

Figure 3A shows a Northern analysis autoradiogram of RNA extracted from the leaves of a brassinosteroid non-sensitive mutant (Akitakomachi) and RNA extracted from various organs of a wild type rice plant (Nipponbare).

Figure 3B shows a Northern analysis autoradiogram of RNA extracted from brassinosteroid non-sensitive mutants and RNA extracted from wild type rice plants. The left-hand side of Figure 3B shows a comparison between wild types and mutants obtained by using a 5' probe. The right-hand side of Figure 3B shows a comparison between wild types and mutants obtained by using a 3' probe.

Figure 4 shows an amino acid sequence of the novel rice gene which controls a physiological reaction system induced by brassinosteroid hormone, together with characteristic sequences found therein (where nuclear localization signals and an ATP/GTP binding motif can be observed).

Figure 5A shows a brassinosteroid leaf blade bending experimentation using a mutated line (A0369) derived

from Akitakomachi. The left-hand side shows results of wild type plants, whereas the right-hand side shows results of mutants.

5 Figure 5B shows a brassinosteroid leaf blade bending experimentation using a mutated line (NC6148) derived from Nipponbare. The left-hand side shows results of wild type plants, whereas the right-hand side shows results of mutants.

10 DESCRIPTION OF THE PREFERRED EMBODIMENTS

 The present invention provides a method for producing an improved plant, the method utilizing a novel
15 plant gene which can be provided by using Tos17.

 According to the present invention, there is provided a polynucleotide encoding a plant gene capable of controlling various effects in which brassinosteroid
20 hormone is involved. As used herein, the term "capable of controlling various effects" means the ability to control various effects in plants in which brassinosteroid hormone is involved, e.g., growth promotion, yield increase, quality improvement, maturation enhancement, and tolerance against
25 biotic and abiotic stresses, including dwarfism, upright form, and malformation of grain hulls, thereby providing a number of agriculturally useful effects as are attained by treatments with brassinosteroid hormone agricultural chemicals. The term "plants" encompasses both
30 monocotyledons and dicotyledons.

 A polynucleotide encoding a plant gene capable of controlling a signal transduction system for

brassinosteroid hormone according to the present invention is, for example, a polynucleotide encoding an amino acid sequence from Met at position 1 to Arg at position 1057 of SEQ ID NO: 2 in the SEQUENCE LISTING, including any
5 polynucleotide encoding an amino acid sequence in which one or more amino acids are deleted, substituted or added to the aforementioned amino acid sequence.

10 A polynucleotide encoding a plant gene capable of controlling various effects in which brassinosteroid hormone is involved encompasses any polynucleotides which have at least about 80% sequence homology, preferably at least about 85% sequence homology, and more preferably at least about 90% sequence homology, still more preferably
15 at least about 95% sequence homology, and most preferably at least about 99% sequence homology, with an amino acid sequence from Met at position 1 to Arg at position 1057 of SEQ ID NO: 2 in the SEQUENCE LISTING, so long as they are capable of controlling various effects in plants in which
20 brassinosteroid hormone is involved. The term "sequence homology" indicates a degree of identity between two polynucleotide sequences to be compared with each other. The rate (%) of sequence homology between two polynucleotide sequences for comparison is calculated by, after optimally
25 aligning the two polynucleotide sequences for comparison, obtaining a matched position number indicating the number of positions at which identical ("matched") nucleic acid bases (e.g., A, T, C, G, U, or I) are present in both sequences, dividing the matched position number by total number of bases
30 in the polynucleotide sequences for comparison, and multiplying the quotient by 100. The sequence homology can be calculated by using the following sequencing tools, for example: a Unix base program designated GCG Wisconsin

Package (Program Manual for the Wisconsin Package, Version 8, September 1994, Genetics Computer Group, 575 Science Drive Madison, Wisconsin, USA 53711; Rice, P. (1996) Program Manual for EGCG Package, Peter Rice, The Sanger Centre, Hinxton Hall, Cambridge, CB10 1RQ, England), and the ExPASy World Wide Web molecular biology server (Geneva University Hospital and University of Geneva, Geneva, Switzerland).

Cells into which genes have been introduced are first selected based on drug resistance, e.g., hygromycin resistance, and then regenerated into plant bodies by using usual methods.

The terminology and laboratory procedures described throughout the present specification are directed to those which are well-known and commonly employed in the art. Standard techniques may be used for recombination methods, polynucleotide synthesis, microorganisms culturing, and transformation (e.g., electroporation). Such techniques and procedures are generally known from various standard textbooks available in the field or by way of the present specification (including a generally-referenced textbook by Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd edition (1989) Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.). Such literature is incorporated herein by reference.

The polynucleotide according to the present invention can be obtained by using the method described herein, for example. However, the polynucleotide according to the present invention may also be obtained by any chemical synthesis process based on the sequence disclosed herein. For example, the polynucleotide according to the present

invention may be synthesized by using a polynucleotide synthesizer available from Applied Bio Systems in accordance with the instructions provided by the manufacturer.

5 Methods of PCR amplification are well-known in the art (PCR Technology: Principles and Applications for DNA Amplification, ed. HA Erlich, Freeman Press, NewYork, NY (1992); PCR Protocols: A Guide to Methods and Applications, Innis, Gelfland, Snisky, and White, Academic Press, San
10 Diego, CA(1990); Mattila et al. (1991) Nucleic Acids Res. 19: 4967; Eckert, K.A. and Kunkel, T. A. (1991) PCR Methods and Applications 1: 17; PCR, McPherson, Quirk, and Taylor, IRL Press, Oxford). Such literature is incorporated herein by reference.

15

(Examples)

 Hereinafter, the present invention will be described by way of examples which are of illustrative but non-limiting nature.

20

(Example 1: Activation of Tos17 via culture)

 Using fully ripened seeds of Nipponbare and Akitakomachi, which are varieties of Japonica subspecies, induction of calli and cell suspension culture were carried
25 out as described earlier (Hirochika et al., 1996, supra). The activation of Tos17 which was used for gene destruction was carried out following the method of Ohtsuki (1990) (rice protoplast culture system, Food and Agricultural Research Development Association). In summary, fully ripened seeds
30 of rice were cultured in an MS medium having 2,4-dichlorophenoxyacetic acid (2,4-D) added thereto (2 mg/ml) (Ohtsuki (1990), supra) (25°C, 1 month), to induce callus formation. The resultant calluses were cultured for

5 months in an N6 liquid medium having 2,4-D added thereto (Ohtsuki (1990), supra), and thereafter placed on a redifferentiation medium (Ohtsuki (1990), supra), whereby redifferentiated rice plants were obtained (first generation (R1) plants).

(Example 2: Isolation of sequences adjoining Tos17)

Utilizing each of the regenerated R1 rice plants obtained according to Example 1 as a first strain, about 1000 R1 seeds were collected from each strain and grown on a paddy field to obtain second generation (R2) plants, which were subjected to a morphological analysis. As a result of observing the phenotypes of the respective plant bodies in the R2 group, it was learned that about 1/4 of the R2 group of an Akitakomachi strain A0369 exhibit the "dwarfism, upright form, and malformation of grain hulls" phenotype (Figures 1A and 1B). In the regenerated group of Akitakomachi, dwarfism, upright form, and malformation of grain hulls were observed for brassinosteroid insensitive mutants (Figure 1A, left, and Figure 1B, left), as compared with the wild type (Figure 1A, right, and Figure 1B, right). The isolation of adjacent sequences of transposed Tos17, which is co-segregating with the phenotypes, was carried out by an IPCR method (Ochman et al., Genetics Nov; 120(3): 621-3(1988) and Triglia et al., Nucleic Acids Res Aug 25; (16): 8186(1988)). The total DNA of A0369 was digested with XbaI, and a ligation process was performed in a large quantity of solution, thereby obtaining self-ligated circular molecules. In the self-ligated circular molecules, the adjacent sequences are flanking the internal sequence of Tos17. As a result, amplification was successfully carried out by usual PCR methods using an outward primer pair (T17TAIL3: GAGAGCATCATCGGTTACATCTTCTC; T17-1950R:

TCTAGCAGTCTCAATGATGTGGCG) based on the known sequence of Tos17.

(Example 3: Search for alleles)

5 Using the sequence obtained according to Example 2,
lineage in which Tos17 had been inserted at a different site
in the same gene was selected by PCR from the regenerated
rice group of Nipponbare. As a result, a line (NC6148) which
similarly exhibited dwarfism, upright form, and
10 malformation of grain hulls were observed for
brassinosteroid was selected. That is, in the regenerated
rice group of Nipponbare, as well, dwarfism, upright form,
and malformation of grain hulls were observed for
brassinosteroid insensitive mutants (Figure 2A, left, and
15 Figure 2B, left), relative to the wild type (Figure 2A,
right, and Figure 2B, right). It was concluded that these
common mutations were results of the same gene having been
destroyed.

20 (Example 4: Expression and analysis of the causative gene)

From the group of R2 rice plants (selfed progeny from
the A0369 and NC6148 strains) obtained according to
Examples 2 and 3, individuals exhibiting mutation were
identified from normal individuals. RNA was prepared from
25 both groups of individuals, and the expression specificity
was analyzed through Northern analysis. After agarose
electrophoresis, the RNA obtained from individuals
exhibiting mutation and the RNA obtained from normal
individuals were allowed to adsorb to nylon membranes. DNA
30 fragments which were obtained by amplifying via PCR a
sequence (positions 5775-6638 of the genomic sequence) on
the 5' side and a sequence (positions 8175-8765 of the
genomic sequence) on the 3' side of the Tos17 insertion site

in both mutated lines were labeled with ^{32}P -dCTP. By using these as probes, expression specificity was analyzed through Northern analysis (Figures 3A and 3B). As seen from the Northern analysis autoradiogram shown in Figure 3A, a band (about 4.3 kb) indicated by an arrow was confirmed to be expressed in all observed organs of the wild type. However, in the mutants, transcripts of abnormal sizes were observed due to the insertion of Tos17, indicating that the natural function of the wild type is lost (Figure 3B).

(Example 5: Structural analysis of the causative gene)

Using the sequence obtained according to Example 2 as a probe, the corresponding cDNA and genomic clone were obtained from a cDNA library and a genomic library. Their structures are shown in SEQ ID Nos: 1 and 3. It was learned that this gene includes 6 exons and 5 introns, encoding 1057 amino acids, and that Tos17 had been inserted at the 4th and 5th exons in two mutants, respectively. Moreover, motif search results suggested the presence of nuclear localization signal 1 (amino acid residues 329-367 of SEQ ID NO: 2, Robbins & Dingwall consensus sequence; a search result by PSORT program (<http://psort.ims.u-tokyo.ac.jp/>)) and nuclear localization signal 2 (amino acid residues 457-460, 595-600 of SEQ ID NO: 2, 4 amino acid nuclear localization pattern signal; a search result by PSORT program (<http://psort.ims.u-tokyo.ac.jp/>)) as well as the presence of an ATP/GTP binding domain (amino acid residues 526-533 of SEQ ID NO: 2; a search result by a motif search service on Genomenet(<http://www.genome.ad.jp/>)). Thus, the possibility of this gene being involved in signal transduction was suggested (Figure 4).

(Example 6: brassinosteroid sensitivity evaluation)

5 The present gene was deduced to be a factor involved
in the signal transduction system for plant hormones, taking
note of the facts that the present gene was expressed in
all plant bodies and that pleiotropic influences resulted
from destroying this gene, as well as the possibility that
the gene might be a factor involved in the signal
transduction system. Presuming that the signal
transduction system is that for brassinosteroid hormone in
view of the resultant upright form, the inventors performed
10 a leaf blade bend response test as a brassinosteroid response
test, by using brassinolide, which is one kind of
brassinosteroid hormone. The second leaf of rice which was
allowed to germinate in the dark was cut off, and immersed
in a 1 ng/ml of brassinosteroid solution for 48 hours. The
15 wild type individuals having the wild type genes showed
bending of the leaf blades and leaf sheath junctions
(left-hand side in Figures 5A and 5B), showing response to
brassinolide, whereas mutant individuals showed little
bending thereof (right-hand side in Figures 5A and 5B),
20 indicating that the destruction of the present gene resulted
in the loss of response to brassinosteroid. From the above
results, it was revealed that the present gene is a gene
involved in the signal transduction system for
brassinosteroid hormone.

25 The above examples are illustrative, and by no means
limiting, of various aspects of the present invention and
the manners in which the oligonucleotide according to the
present invention can be made and utilized.

30 Thus, according to the present invention, a novel
polynucleotide is provided which is capable of controlling
various effects in which brassinosteroid hormone is involved,

the polynucleotide being of use in plant breeding. By introducing the present polynucleotide into plants and artificially controlling various effects in which brassinosteroid hormone is involved, it is expected that effects such as growth promotion, yield increase, quality improvement, maturation enhancement, and tolerance against biotic and abiotic stresses can be controlled, thereby providing a number of agriculturally useful effects as are attained by treatments with brassinosteroid hormone agricultural chemicals.

Various other modifications will be apparent to and can be readily made by those skilled in the art without departing from the scope and spirit of this invention. Accordingly, it is not intended that the scope of the claims appended hereto be limited to the description as set forth herein, but rather that the claims be broadly construed.

SEQUENCE LISTING

<110> National Institute of Agrobiological Resources, Ministry of Agriculture,
Forestry and Fisheries

5 Japan Science and Technology Corporation

<120> A novel gene involved in brassinosteroid responses

<130>

10

<140> JP

<141> 2000-05-22

<160> 3

15

<170> PatentIn Ver. 2.1

<210> 1

<211> 4310

20

<212> DNA

<213> Oryza sativa

<220>

<221> CDS

25

<222> (655).. (3828)

<400> 1

ctcttctcca ctccaaatcc cttcttacc ctttccctc cccccgcagc ttctcttct 60

30

cctgcagtac tcgccgccac caccaccgcg ccgccgccgc cggccgcgtt ccgagaccca 120

ctcgatcgga atccaccgcg gcgcgcccgc gcgccctgcgt cctcttctt ccccgaggagc 180

[illegible]

5

10

15

20

25

30

gtg caa gaa ggc aat gac tcc aaa tgc aat gcg cct tct ggc aag aat 1233
Val Gln Glu Gly Asn Asp Ser Lys Cys Asn Ala Pro Ser Gly Lys Asn
180 185 190

195

cca tgt gaa gaa gtt gtt ctc aaa aga agc tcc aaa tct aag agg aag 1665

ctt atg aac tgg ctg aat gga aaa aag aaa aga act gga agt gtg cat 2049
Leu Met Asn Trp Leu Asn Gly Lys Lys Lys Arg Thr Gly Ser Val His

	450	455	460	465	
	cac aca gtt gct cat cca gct ggg aat ttg agc aac aaa aaa gtg aca				2097
	His Thr Val Ala His Pro Ala Gly Asn Leu Ser Asn Lys Lys Val Thr				
5		470	475	480	
	ccc act gcg agt act cag cat gat gat gag aat gat act gaa aat ggt				2145
	Pro Thr Ala Ser Thr Gln His Asp Asp Glu Asn Asp Thr Glu Asn Gly				
		485	490	495	
10					
	ctt gac aca aat atg cat aag aca gat gtc tgt cag cat gta tca gaa				2193
	Leu Asp Thr Asn Met His Lys Thr Asp Val Cys Gln His Val Ser Glu				
		500	505	510	
15					
	atc tcc aca cag agg tgc tca tca aag ggg aaa aca gcg ggt ttg agt				2241
	Ile Ser Thr Gln Arg Cys Ser Ser Lys Gly Lys Thr Ala Gly Leu Ser				
		515	520	525	
	aag ggg aaa aca cat tca gct gct agt acc aaa tat ggt ggt gaa agc				2289
20	Lys Gly Lys Thr His Ser Ala Ala Ser Thr Lys Tyr Gly Gly Glu Ser				
		530	535	540	545
	acc aga aat ggt cag aac ata cat gta ctc agc gca gaa gat caa tgc				2337
	Thr Arg Asn Gly Gln Asn Ile His Val Leu Ser Ala Glu Asp Gln Cys				
25		550	555	560	
	cag atg gaa acc gaa aac tct gtt ctg agt cac tcg gca aag gtt tct				2385
	Gln Met Glu Thr Glu Asn Ser Val Leu Ser His Ser Ala Lys Val Ser				
		565	570	575	
30					
	cca gct gag cat gat atc caa att atg tct gac ctt cat gag cag agt				2433
	Pro Ala Glu His Asp Ile Gln Ile Met Ser Asp Leu His Glu Gln Ser				
		580	585	590	

5

10

15

20

25

30

cag ggt cat ttg gca ttg acc aca caa gag tct cca cat cct cag aac 2769
Gln Gly His Leu Ala Leu Thr Thr Gln Glu Ser Pro His Pro Gln Asn
690 695 700 705

ttt cag tct act cag gaa cag cag aca cat ttg cgg atg gaa gaa atg 2817
Phe Gln Ser Thr Gln Glu Gln Gln Thr His Leu Arg Met Glu Glu Met
710 715 720

	980	985	990	
	cag ttg ggt tct gtt agc tcc agt gca gat ttt tta tca gcg agg aac			3681
	Gln Leu Gly Ser Val Ser Ser Ser Ala Asp Phe Leu Ser Ala Arg Asn			
5	995	1000	1005	
	agc ata gct caa tct tgg acc aga ggc aag ggt aaa atg gtt cat ccc			3729
	Ser Ile Ala Gln Ser Trp Thr Arg Gly Lys Gly Lys Met Val His Pro			
	1010	1015	1020	1025
10	ttg gat cgg ttt gtg aga cag gat atc tgt ata act aac aag aac cca			3777
	Leu Asp Arg Phe Val Arg Gln Asp Ile Cys Ile Thr Asn Lys Asn Pro			
	1030	1035	1040	
15	gct gat ttt act aca atc agt aac gat aac gag tat atg gat tac cgc			3825
	Ala Asp Phe Thr Thr Ile Ser Asn Asp Asn Glu Tyr Met Asp Tyr Arg			
	1045	1050	1055	
	tga agcagaaagt ggtgtgcata attcctgaac atttacaatc atacatttca			3878
20				
	tcctttatggc gccaaatagt catactgtaa gaggagggct ttgctggatc tgctgtgaagg			3938
	cttcttgtaa gtgtgtggatg ccccatcttc tggatgggaa cctgccagac agtgaacaag			3998
25	ggctttgcaa ggtgcagcat ccggtttttg ttttgccagt ccaagaaacg tcctcctgtt			4058
	actttgtagt tgtactcata ctagtgcgct tgtttgtaca aggagaaatg tgtaaccttg			4118
30	ttgaaaaaat gtctcccca ttttgtaatt accataagga ggtttatagt gtgtgagct			4178
	gtgtgtgact gacggcgaga aatggttttg tgggtgttaa ggttgaaacg actagctctc			4238

ggagaattta ca 4310

<210> 2
<211> 1057
<212> PRT
<213> *Oryza sativa*

<400> 2																		
Met	Glu	Ile	Val	Ala	Val	Asp	Gln	Glu	Gly	Ala	Arg	Val	Val	Gly	Thr			
1				5					10					15				
Asn	Cys	Met	Leu	Ala	Arg	Gly	Gly	Thr	Gly	Ala	Val	Ala	Pro	Val	Leu			
			20					25					30					
Glu	Leu	Thr	Ala	Thr	Pro	Arg	Gln	Asp	Ala	Ala	Ala	Glu	Ala	Gly	Val			
		35					40					45						
Asp	Glu	Pro	Ala	Gln	His	Gln	Cys	Glu	His	Phe	Ser	Ile	Arg	Gly	Tyr			
	50					55					60							
Val	Ala	Leu	Leu	Gln	Lys	Lys	Asp	Pro	Lys	Phe	Cys	Ser	Leu	Ser	Arg			
65				70						75					80			
Ile	Phe	His	Asp	Gln	Lys	Lys	Cys	Asp	Glu	His	Lys	Ala	Ser	Ser	Ser			
			85						90					95				
Pro	Phe	Ser	Val	Ala	Lys	Phe	Arg	Arg	Trp	Asp	Cys	Ser	Lys	Cys	Leu			
			100					105					110					
Asp	Lys	Leu	Lys	Thr	Ser	Asp	Asn	Gly	Thr	Ala	Pro	Arg	Thr	Leu	Pro			
		115					120					125						
Ala	Lys	Gln	Asn	Gly	Thr	Ser	Asp	Gly	Cys	Ser	Ile	Thr	Phe	Val	Arg			
	130					135					140							
Ser	Thr	Phe	Val	Pro	Ala	Ser	Val	Gly	Ser	Gln	Lys	Val	Ser	Pro	Ser			
145				150					155					160				
Thr	Gln	Ser	Ser	Gln	Gly	Lys	Asn	Ala	Asp	Arg	Ser	Thr	Leu	Pro	Lys			
			165					170					175					

Ser Val Gln Glu Gly Asn Asp Ser Lys Cys Asn Ala Pro Ser Gly Lys
 180 185 190
 Asn Gly Ala Ala Glu Ala Asn Thr Asp Ser Pro Met Lys Asp Leu Gln
 195 200 205
 5 Gly Pro Ala Gln Asn Tyr Asp Val Ala Ala Asn Val Ser Glu Asp Asn
 210 215 220
 Thr Ser Val Asp Val Gly Ala Leu Pro Glu Val Pro Gln Ile Thr Trp
 225 230 235 240
 His Ile Glu Val Asn Gly Ala Asp Gln Pro Pro Ser Thr Pro Lys Leu
 10 245 250 255
 Ser Glu Val Val Leu Lys Arg Asn Glu Asp Glu Asn Gly Lys Thr Glu
 260 265 270
 Glu Thr Leu Val Ala Glu Gln Cys Asn Leu Thr Lys Asp Pro Asn Pro
 275 280 285
 15 Met Ser Gly Lys Glu Arg Asp Gln Val Ala Glu Gln Cys Asn Leu Thr
 290 295 300
 Lys Asp Pro Lys Pro Val Ser Gly Gln Lys Cys Glu Gln Ile Cys Asn
 305 310 315 320
 Glu Pro Cys Glu Glu Val Val Leu Lys Arg Ser Ser Lys Ser Lys Arg
 20 325 330 335
 Lys Thr Asp Lys Lys Leu Met Lys Lys Gln Gln His Ser Lys Lys Arg
 340 345 350
 Thr Ala Gln Ala Asp Val Ser Asp Ala Lys Leu Cys Arg Arg Lys Pro
 355 360 365
 25 Lys Lys Val Arg Leu Leu Ser Glu Ile Ile Asn Ala Asn Gln Val Glu
 370 375 380
 Asp Ser Arg Ser Asp Glu Val His Arg Glu Asn Ala Ala Asp Pro Cys
 385 390 395 400
 Glu Asp Asp Arg Ser Thr Ile Pro Val Pro Met Glu Val Ser Met Asp
 30 405 410 415
 Ile Pro Val Ser Asn His Thr Val Gly Glu Asp Gly Leu Lys Ser Ser
 420 425 430
 Lys Asn Lys Thr Lys Arg Lys Tyr Ser Asp Val Val Asp Asp Gly Ser

002214-14300

965 970 975
 His Phe Glu Ala Leu Asn Ser Gly Met Phe Ser Ala Lys Trp Asn Ala
 980 985 990
 Leu Gln Leu Gly Ser Val Ser Ser Ser Ala Asp Phe Leu Ser Ala Arg
 5 995 1000 1005
 Asn Ser Ile Ala Gln Ser Trp Thr Arg Gly Lys Gly Lys Met Val His
 1010 1015 1020
 Pro Leu Asp Arg Phe Val Arg Gln Asp Ile Cys Ile Thr Asn Lys Asn
 1025 1030 1035 1040
 10 Pro Ala Asp Phe Thr Thr Ile Ser Asn Asp Asn Glu Tyr Met Asp Tyr
 1045 1050 1055
 Arg

 <210> 3
 15 <211> 9455
 <212> DNA
 <213> *Oryza sativa*

 <400> 3
 20 gatctatcgt tgattggatt tcgctgggct agctaggttag acaagggttc aaaatgtgac 60
 gaacattatt gagctaataa ataacgagaa accaacttgg tatataggtc aatttcaaaa 120
 gaaacaagct gacaaaattc gtccaatttc actagttttt gtcagtaatt gaatggcaat 180
 catggttatac gacaaaaccg cttaggagtg ctatttgggtg atggaatggt ttataaaaact 240
 ttggaccgga gtagcagtac aatggcttgt ctgaacaggc taggtagcat agtaggtcct 300
 25 tttagccttgg ttgcactgtt ctgtcggcct ataggggaat ctattggcctt aatggaaggg 360
 aaaatagtgt gactagcatc atcgatttgc ttgctatcat gtttgagcat cattgacatg 420
 tgggtgtcga tctaggagac tatgaatcta gcgaatcctc actagtatgc acatgcacaa 480
 cgacatcatg cagctataatg tacaacaatg taggaatggt agctctactc ggattggttg 540
 ccgcattcct acattgctag aaaaacacaa acacacacac acacacacac aaacaacat 600
 30 aggacgtcgc aacacctctt ccctggctac cactatcgcc aaatgccaag catggttgtg 660
 ttgacctccc tccttagctg ccactgtgtc cacttcaaaa tccataacca cgacatcttc 720
 ccttcttcta caattgttac tactgttctt gaagagggtc ttgtcctacc gacttactaa 780
 agaccaacca ttggtcagca gagcatctcg agacttctat accaaagtaa cattgagagg 840

0022141300

	tctcttctcc	actccaaatc	ccttcttacc	ctatttccct	ccccccgcag	cttctcttcc	2880
	tccitgcagta	ctcgcgcgcca	ccaccaccgc	gccgcgcgccg	cgggcgcggt	tccgagacc	2940
	actcgaicgg	aatccaccgc	ggcgcgcgccg	cgcgccitgcg	tcctcttct	tccccgggag	3000
	ccgaccgacc	acggcgacca	gtcgaatctc	ctctccgggc	gccaaccgcg	tcttagcttc	3060
5	atcgaatcca	ccgccccacc	ccgcattctc	tcctctctct	ccgacgacga	cgactactac	3120
	tagtcttctc	caataagccc	ccctcccgct	ccccccgcct	gaagaagaag	cagcagctag	3180
	ctccggggag	aggctgcacg	cgcgccgggt	agatcgcgcc	ccgccccgcc	tgcgtcgcgg	3240
	ctgtcggagc	aaacgcaaac	ccccaggta	atcaacgaac	ttttctccg	ccgcaagaac	3300
	agctcccgcg	gggggtttgg	ttttgaccga	tttcttcccc	cctcccccca	aatcgaccca	3360
10	tccaatttgc	cctcgattta	cttccgattt	cccactttt	ttttcttct	ttcgggttgg	3420
	gggggttgcg	ttttggggga	ggagagggtt	tcagctcatc	cgaagcccca	cgttaggttc	3480
	gcccccttfc	cagctgtgcc	cctctctcgg	gcctcgagct	cctcgccctc	atgggaacca	3540
	aagcccttat	atttcatgtc	gcggaagaaa	aaaaaatccc	gtcttttgcg	gggatcctcg	3600
	cggctacgta	cgagccctag	ttaccgcgcg	gattttagtt	acggcgggtt	atgcggcccc	3660
15	tccctctagg	ttttagatct	acccatctct	ctctctctct	ctctctctct	ctctgtgcat	3720
	gcattgtgtc	atcttagcta	tacctgtatt	atttgggaagg	ttaattatgg	ttgtgtatat	3780
	gtggcgcggt	aattaattag	tttaattcgc	acccccctct	tctttgttta	tctaggtttt	3840
	gggggaattt	atttcttgc	ataattttgc	ccgctcgaat	ttctggtgct	cttatatttc	3900
	atgagctgat	tgaagtggat	atatattgtg	cgctgcgtgcg	tgctattgct	acatcggctt	3960
20	gacttcttct	tgcctactac	ttcatttaatt	tgtttcttct	ggtttctgtt	tcaggttgtt	4020
	ctagcgtgtg	cagcggctag	ctgattgatt	gtcttctgtg	atataaccag	agctcgtgtt	4080
	ttgtggtttg	tggtttgtgg	tttgtgttgg	gattgttgat	gtgctaattc	gcggcgttac	4140
	aagatcactg	ctggattgat	attgagttgt	gcctcggctg	tgctagctgt	gtgttgattc	4200
	tctctctcgc	gtgggtgatcg	atatggagat	tgttcgagta	gatcaggagg	gagctcgtgt	4260
25	tgttgggacg	aactgtatgc	ttgctcgtgg	tggaaactgg	gctgtagcgc	cagtgttggg	4320
	gctgacagcg	acgcctcgtc	aggatgcagc	cgctgaagct	gggttagacg	aaccggcaca	4380
	acaccaatgc	gagcatttct	ccataaggta	atcattttct	gtatttccaa	ttccagtatc	4440
	gcgttgttga	tgaataatga	atcgccatgt	catgccatat	tgcactgttt	gatggaagag	4500
	tatgattgat	cgtgggtttt	gcacagtttg	ctgttgggac	ttatatggtc	atctgttttg	4560
30	tacgatcgta	tacactgggt	cgacatgctt	atgactttgg	ttcgatttag	gaagtcaata	4620
	catccactac	tagctctata	tctagccatg	tgaactcatt	tatgccatag	cacagctagc	4680
	aggctagcag	caaaaaatat	atataatat	tgcataatat	ttgggtgttc	atgtatcttt	4740
	atactctacg	tacatccatt	aatacttcca	atgtatgaat	ctgagcacat	gattgtgagt	4800

gclacacata tgcattgtctg tatgtgtgtt cattagggtt ttgatcatat ttgttttgtt 4860
 tgggggtgcgc atgcatittat tcaggccatg ctgtaggctg tagctagata ttgtgttttg 4920
 tatattattt ctgttgaaca agctgattac taatgaaatt aacctttttg gggtagactc 4980
 atatatggg ccctacattt ttgtaatcat ttttcctttg tgcigagggt cagcataaaa 5040
 5 cttttttatc alaagcatgt ttacatccta ggagattctt agaactgatg gtttcttcat 5100
 atttgcatta tgtttgattt gatagtcctt tattatttta agccttttca attgttttaga 5160
 gattctagag atgatatata tcaaccatag acttgtcagc ttttggttta atacitttcta 5220
 gaactaatta gattattatt ttgttagttt atcctgtcat gctatttgta ttatctttga 5280
 attcaaactg caatacttag attatcttga aggtcctctt tttctggact gtacaagcta 5340
 10 tgtatgaaat gcctacctcc cagcatcctt tagattatgt agggcctttt ctgagtttat 5400
 cagttgtata ttgactgaag cagcgaatgt gctatatata tgtgccatgc atctttataa 5460
 tgataatctt atttttcttg taccagaggg tatgttgctc ttcttcagaa gaaggatcca 5520
 aaattctgct ctctatctcg gatittccat gaccagaaaa aatgtgatga acacaaagct 5580
 agttcaagcc catittctgt agcaaagttt cgacgatggg attgctcgaa gtgcttggat 5640
 15 aagttgaaaa cttcagataa tggaacagca ccaagaactc ttcccgcaa gcagaatggc 5700
 acaagtgatg gttgctccat cacatttgtt cggagcactt ttgtgcctgc tagtgttgg 5760
 tcccaaaaag tgtctcctag cacacaatca tctcaaggga agaattgctga tagatcaact 5820
 ctccaaaaga gtgtgcaaga aggcaatgac tccaaatgca atgcgccttc tggcaagaat 5880
 ggagctgctg aggccaatac tgattcacca atgaaaggta tggtagatgt agagcctttc 5940
 20 aaattcctaa gtaggatttt atttaaggta tagaataaac taatgtttgt gtgattttct 6000
 cagatttgca agggccagcc caaaattatg atgtggcagc aaatgtctct gaggacaaca 6060
 ctctgttga tgttggggct ttacctgaag tccccagat tacatggcac atagaagtaa 6120
 atgggtgcaga tcaacctcca tccactcaa aactttctga agtggctctc aaaagaaatg 6180
 aagatgaaaa tggaaaaact gaagagactc ttgttgctga gcagtgaat ttgaccaaag 6240
 25 atcctaacc aatgtctgga aaggaacgtg atcaggttgc tgagcagtgc aatttgacca 6300
 aagatccgaa accagtgtct gggcagaaat gtgagcagat ctgcaatgag ccatgtgaag 6360
 aagttgttct caaaagaagc tccaaatcta agaggaagac ggataagaag ttgatgaaga 6420
 agcagcagca cagcaagaaa cgcactgccc aggcigatgt ttcatatgca aagctttgtc 6480
 ggagaaagcc aaaaaagggt cggcttctat cagaaattat aaatgctaac caggttgagg 6540
 30 attctagaag tgacgaagtt catcgtgaaa atgccgtga tccctgtgag gatgatagaa 6600
 gtaccatccc ggtcccgatg gaagtaagca tggatatcc tttagcaac catacagtgg 6660
 gagaagatgg gttaaaatca agtaagaaca agacaaaacg caaatactct gatgtttag 6720
 atgatggatc atcattatg aactggctga atggaaaaa gaaaagaact ggaagtgtgc 6780

0022144300

atcacacagt tgctcatcca gctgggaatt tgagcaacaa aaaagtgaca cccactgcga 6840
 gtactcagca tgatgatgag aatgatactg aaaaatggctt tgacacaaat atgcataaga 6900
 cagatgtctg tcagcatgta tcagaaatct ccacacagag gtgctcatca aaggggaaaa 6960
 cagcgggitt gagtaagggg aaaacacatt cagctgctag taccaaataat ggtggtagaa 7020
 5 gcaccagaaa tggtcagaac atacatgtac tcagcgcaga agatcaatgc cagatggaaa 7080
 ccgaaaactc tgttctgagt cactcggcaa aggtacgaat tttgtgaatc atgaggaatt 7140
 tttgcttttt aaattgactg aatcaacatt tatctgtatg aaggaataat attggtgcat 7200
 aacaatgita agaaatatgc atacaatgtt tatttataatg ctttccactg ttcttcttta 7260
 cttatgtttt gatactcttt ttgtgtgtgc gtgcatgtgt gcatgtgtgt gtgtgtgtgt 7320
 10 gtgtgcgcgc gtgtgtgtgc acgtgcgtgg cgcaatattc ttttttagac tcatattata 7380
 gtgattgtaa tggactgaca ttttctcat ttctcatctc aggtttctcc agctgagcat 7440
 gatatccaaa ttatgtctga cttcatgag cagagcttac ccaagaagaa aaagaagcaa 7500
 aaacttgaag tgactcgtga aaaacagacc atgatagatg acatcccat ggatattgtt 7560
 gaactgctag ctaaaaacca gcatgagagg cagcttatga ctgagactga ttgttctgac 7620
 15 atcaaccgtt ttcaatccaa gacaactgct gatgatgatt gtgtaatagt agctgccaag 7680
 gatggttcag attatgcatac aagtgtgttt gacactaatt cccaacagaa gtccttggca 7740
 tcccaaagta cacagaagga gttacagggt catttggcat tgaccacaca agagtctcca 7800
 catcttcaga acttcagtc tactcaggaa cagcagacac atttgcggat ggaagaaatg 7860
 gtcactattg ctgcaagctc accactattt tcacatcatg atgatcagta tattgctgaa 7920
 20 gcaccaactg aacattgggg ccgtaaggac gcaaagaagc taacgtggga gcaatttaag 7980
 gccactacaa gaaattctcc agcagcaaca tgtgggtgctc aatttagacc tggatatcaa 8040
 gcagttgact tgacttctac tcatgtcatg ggaacttcca gcaattatgc atctcgccaa 8100
 ccagtaattg cgccactgga ccgctatgct gaaagagcgg ttaaccaggt ccatgcaaga 8160
 aattttccaa gcacaatagc aaccatggaa gcgagtaagt tatgtgatcg gagaaatgct 8220
 25 ggacaagtag tcttgatcc taaagaatcc atgcctgcga cgcatcttct gagaatgatg 8280
 gatccatcaa cattagcaag cttccccaac tatggaactt ctagcaggaa ccagatggag 8340
 tctcaacttc ataattctca gtatgcacat aatcagtaca aaggatcaac cagcacatca 8400
 tatggcagta acctgaatgg aaagattcca ttgacattcg aagacttatac acggcatcag 8460
 ctgcatgatc tgcacagacc tttagccca catcctagag ttgggtgtgt tggctccttg 8520
 30 ctgcagaagg aaattgcaaa ctggtcggag aactgtggca cacaatctgg ttataagtta 8580
 ggagtgtaaa caggaataac atcgcatcag atgaacagaa aggaacattt tgaagccctg 8640
 aattctggaa tgttttcagc aaaatggaat gcattgcagt tgggttctgt tagctccagt 8700
 gcagattttt tatcagcgag gaacagcata gctcaatctt ggaccagagg caagggtaaa 8760

0022114-11200

atggttcac ccttggatcg gtttgtgaga caggatatct gtataactaa caagaaccca 8820
gctgatttta ctacaatcag taacgataac gagtatatgg attaccgctg aagcagaaag 8880
tgggtgtgcat aattcctgaa catttacaat catacatitc atctttatgg cgccaaatag 8940
tcatactgta agaggagggc ttigtctggat ctgctgtaag gtaagttgaa ctttttcttc 9000
5 tlgcaagttt atcagtttaa gaaaaaagaa tgattactta tggtagcaag gatggttctt 9060
gcaggcttct tgtaagttgt ggatgcccc ttttctggat gggaacctgc cagacagtga 9120
acaagggctt tgcaaggtgc agcatccggt ttttgttttg ccagtccaag aaacgtcctc 9180
ctgttacttt gtagttgtac tcatactagt gcgcttgitt gtacaaggag aaatgtgtaa 9240
ccttgttgaa aaaatgtctc cccattttg taattacat aaggaggttt atagtgttgt 9300
10 gagctgtgtg tgactgacgg cgagaaaagg ttttgtcggg gtttaaggttg aaacgactag 9360
ctctcgttat caatgtgttg taaacttcta gattgatgtg ttaccttact ctgaagtca 9420
acaccggaga atttacagta cttttttgcc gtgcc 9455

1. A polynucleotide encoding a plant gene capable of controlling a signal transduction system for brassinosteroid hormone, the polynucleotide encoding an amino acid sequence from Met at position 1 to Arg at position 1057 of SEQ ID NO: 2 in the SEQUENCE LISTING, including any polynucleotide encoding an amino acid sequence in which one or more amino acids are deleted, substituted or added to the amino acid sequence.

3. A polynucleotide according to claim 1 as represented by SEQ ID NO: 1 in the SEQUENCE LISTING.

There is provided a polynucleotide encoding a plant gene capable of controlling a signal transduction system for brassinosteroid hormone, the polynucleotide encoding an amino acid sequence from Met at position 1 to Arg at position 1057 of SEQ ID NO: 2 in the SEQUENCE LISTING, including any polynucleotide encoding an amino acid sequence in which one or more amino acids are deleted, substituted or added to the amino acid sequence.

A **Mutant** **Wild Type**
 (A0369 -/-) (Akitakomachi)



B **Mutant** **Wild Type**
 (A0369 -/-) (Akitakomachi)



Fig. 1

[illegible]

Fig. 2

Nipponbare Wild Type		Akitakomachi Wild Type (leaf)	
seedling		seedling	
root		root	
leaf		leaf	
callus		callus	

seedling
ro


root

leaf

callus.

Akitakomachi
Wild Type

Wild Type (leaf)



Wild Type band

Fig. 3A

002277 11112260




Fig. 3B

MEIVAVDQEG ARVVGTCNML ARGGTGAVAP VLELTATPRQ DAAAEAGVDE 50

PAQHQCEHFS IRGYVALLQK KDPKFCSLR IFHDQKKCDE HKASSSPFSV 100

AKFRRWDCSK CLDKLKTSDN GTAPRTLPAK QNGTSDGCSI TFVRSTFVPA 150

SVGSQKVSPS TQSSQGNAD RSTLPKSVQE GNSKCNAPS GKNGAAEANT 200

DSPMKDLQGP AQNYDVAANV SEDNTSVDVG ALPEVPQITW HIEVNGADQP 250

PSTPKLSEVV LKRNEDENGK TEETLVAEQ NLTKDPNPM GKERDQVAEQ 300 Nuclear localization
signal (1)

CNLT KD PKFV SGQKCEQICN EPCEEVLEK SSKSRKRTDK KLMKKQOHSK 350

KRTAQADVSD AKLCRRKPKK VRLLEIINA NQVEDSRSD VARENAADPC 400

EDDRSTIPVP MEVSMDIPVS NHTVGEDGLK SSKNKTTRKY SDVVDGSSL 450 Nuclear localization
signal (2)

MNWLNGKKKR TGSVHTVAM PAGNLSNKKV TPTASTQHDD ENDTENGLDT 500

NMHKTDVCQH VSEISTQRCS SKGKTAGLSK GAVHSAASTK YGGESTRNGQ 550

NIHVLSAEDQ CQMETENSVL SHSAKVSPAE HDIQIMSDLH EQSLPKKKKK 600

QKLEVTREKQ TMIDDIPMDI VELLAKNQE RQMTENDCS DINRIQSKTT 650

ADDDCVIVAA KGDSDYASSV FDTNSQQKSL ASQSTQKELQ GHLALTQES 700

PHPQNFQSTQ EQQTHLRMEE MVTIAASSPL FSHHDDQYIA EAPTEHWGRK 750

DAKKLTWEQF KATTRNSPAA TCGAQFRPGI QAVDLTSTHV MGSSSNYASR 800

QPVIAPLDYR AERAVNQVHA RNFPSTIATM EASKLCDRN AGQVVLYPKE 850

SMPATHLLRM MDPSTLASFP NYGTSSRNQM ESQIHNSQYA HNQYKGSTST 900 ATP/GTP binding motif

SYGSNLNGKI PLTFEDLSRH QLHDLHRPLR PHPRVGVLGS LLQKEIANWS 950


ENCGTQSGYK LGVSTGITSH QMNRKEHFEA LNSGMFSAKW NALQLGSVSS 1000

SADFLSARNS IAQSWTRGKG KMHVPLDRFV RQDICITNKN PADFTTISND 1050

NEYMDYR 1057

Fig. 4

Mutated line (A0369) derived from Akitakomachi




Wild Type

Mutant

Fig. 5A

Mutated line (NC6148) derived from Nipponbare



Wild Type

Mutant

Fig. 5B

DECLARATION AND POWER OF ATTORNEY
FOR PATENT APPLICATION

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name;

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

A NOVEL GENE INVOLVED IN BRASSINOSTEROID RESPONSES
the specification of which

(check ☒ [X] is attached hereto
one)

[] was filed on _____ as
Application Serial No. _____
and was amended on' _____.
(if applicable)

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I do not know and do not believe that the invention was ever patented or described in any printed publication in any country before my or our invention thereof or more than one year prior to this application.

I do not know and do not believe that the invention was in public use or on sale in the United States of America more than one year prior to this application.

I acknowledge the duty to disclose to the United States Patent and Trademark Office all information known by me to be material to patentability as defined in Title 37, Code of Federal Regulations, § 1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, § 119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or

Send correspondence to: James F. Haley, Jr.
FISH & NEAVE
1251 Avenue of the Americas
New York, New York 10020-1104

Direct telephone calls to: James F. Haley, Jr.
(212) 596-9000

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full name of first inventor Hirohiko Hirochika

First Inventor's signature _____ Date _____

Residence Ibaraki, Japan

Citizenship Japan

Post Office Address c/o Ministry of Agriculture, Forestry and Fisheries, National Institute of Agrobiological Resources
2-1-2, Kannondai, Tsukuba-shi, Ibaraki 305-8602 Japan

Full name of second joint inventor Muneo Yamazaki

Second Inventor's signature _____ Date _____

Residence Ibaraki, Japan

Citizenship Japan

Post Office Address c/o Ministry of Agriculture, Forestry and Fisheries, National Institute of Agrobiological Resources
2-1-2, Kannondai, Tsukuba-shi, Ibaraki 305-8602 Japan

